



Viagra® (sildenafil citrate)

USES: Viagra is used to treat male sexual function problems (erection dysfunction).

HOW TO USE THIS MEDICATION: Viagra is taken orally, as needed, between four hours and one-half hour before sexual activity. Taking Viagra about one hour before sexual activity is most effective. Take this medication only as directed (usually once daily as needed). Sildenafil works with any sexual stimulation to help achieve an erection.

SIDE EFFECTS: When taking Viagra, you may suffer from headaches, stomach upset, nasal stuffiness, flushes, diarrhea and dizziness. If any of these side effects persist or become troublesome, consult your doctor immediately.

PRECAUTIONS: Inform your doctor of your complete medical history (particularly if you suffer from any allergies (especially drug allergies). Also inform your doctor if you have any penis conditions (such as fibrosis/scarring). You must tell your doctor if you have sickle cell anemia, blood system cancers (such as leukemia or myeloma), or Peyronie's disease, history of painful/prolonged erection (priapism), eye problems (retina diseases), liver or kidney disease, bleeding disorders or active stomach ulcers, heart diseases, stroke or severe high or low blood pressure.

It is advised to limit alcohol use while taking this medication as it can increase the severity of unwanted side effects. When rising from a seated or lying position, get up slowly as to avoid dizziness and light-headedness. Caution is advised to the elderly, as they may be more sensitive to the side effects of this drug.

DRUG INTERACTIONS: Tell your doctor about any medicines that you use, (including both prescription and non-prescription) particularly if you take any nitrate medications (for example, nitroglycerin or isosorbide dinitrate), nitroprusside (any "nitric oxide donor" medicines), erythromycin, azole antifungals (e.g., itraconazole, ketoconazole), mibefradil, cimetidine, rifamycins (for example, rifampin) or high blood pressure medicines

NOTES: Viagra should not be shared with others, as they may have a problem that cannot effectively be treated using this drug. Use of Viagra does not act as protection against any sexually transmitted diseases (for example, HIV, Hepatitis B, gonorrhea).

MISSED DOSE: Not applicable.

STORAGE: Store Viagra at room temperature. Keep away from moisture and sunlight. Viagra, as with all medications, should be kept out of the reach of children.

Order online at

www.MediPills.com

Copyright© 2002 MediPills.com